



The in vitro antioxidative properties of the essential oils and methanol extracts of *Satureja spicigera* (K. Koch.) Boiss. and *Satureja cuneifolia* ten

Ozgur Eminagaoglu ^a, Bektas Tepe ^{b,*}, Onder Yumrutas ^b, H. Askin Akpulat ^b,
Dimitra Daferera ^c, Moschos Polissiou ^c, Atalay Sokmen ^b

^a Kafkas University, Artvin Forest Faculty, Department of Forest Engineering, 081000 Artvin, Turkey

^b Cumhuriyet University, Faculty of Science and Literature, Department of Biology, 58140 Sivas, Turkey

^c Laboratory of General Chemistry, Agricultural University of Athens, Iera Odos 75, Athens 118 55, Greece

Received 6 June 2005; received in revised form 20 September 2005; accepted 20 September 2005

Abstract

This study was designed to examine the in vitro antioxidant activities of the essential oil and methanol extracts of *Satureja spicigera* and *S. cuneifolia* from Turkish flora. GC and GC/MS analysis of the essential oils resulted in the identification of 40 and 29 compounds, representing the 99.4% and 99.5% of the oils, respectively. Major constituents of the oils were carvacrol (42.5% and 67.1%), γ -terpinene (21.5% and 15.2%) and *p*-cymene (20.9% and 6.7%), respectively. Methanol extracts were also obtained from the aerial parts of the plants. The samples were subjected to a screening for their possible antioxidant activities by using 2,2-diphenyl-1-picrylhydrazyl (DPPH) and β -carotene–linoleic acid assays. In general, samples obtained from *S. cuneifolia* exerted greater antioxidant activities than did those obtained from *S. spicigera*. In the DPPH test system, free radical-scavenging activity of *S. spicigera* oil was determined to be 127 ± 1.63 μ g/ml, whereas IC₅₀ value of *S. cuneifolia* was 89.1 ± 2.29 μ g/ml. In the β -carotene–linoleic acid test system, antioxidant activities of the oil were $81.7 \pm 1.14\%$ and $93.7 \pm 1.83\%$, respectively. Antioxidant activities of the synthetic antioxidant, BHT, ascorbic acid, curcumin and α -tocopherol were also determined in parallel experiments.

© 2005 Elsevier Ltd. All rights reserved.

Keywords: *Satureja spicigera*, *Satureja cuneifolia*; Antioxidant activity; Essential oil; Methanol extract

1. Introduction

The use of spices and herbs as antioxidants in processed foods is a promising alternative to the use of synthetic antioxidants. Numerous reports of antioxidative activity of spices have appeared, strongly inspired by an increasing consumer interest in “natural” food additives. Most investigations have been performed using different model systems, and the spices have been evaluated, either as whole spices or as extract of spices (Madsen & Bertelsen, 1995).

Major reasons for increasing interest in natural antioxidants are: doubts on the safety of use of synthetic substances (butylated hydroxytoluene and butylated hydroxyanisole), the antioxidative efficacy of a variety of phytochemicals, the consensus that foods rich in certain phytochemicals can affect the aetiology, and pathology of chronic diseases and the ageing process and the public conception that natural compounds are innately safer than synthetic compounds and are thus more commercially acceptable (Dorman & Hiltunen, 2004).

Many members of the genus *Satureja* have aromatic and medicinal characteristics. The aerial parts of these species have distinctive tastes and can be added to stuffing, meat pies, and sausages as a seasoning. Fresh sprigs can be

* Corresponding author. Tel.: +90 346 219 10 10x2907; fax: +90 346 219 11 86.

E-mail address: bektastepe@yahoo.com (B. Tepe).

boiled with pulses, such as peas, beans or lentils, for flavouring or, alternatively, they can be used instead of parsley and chervil as a garnish. The leaves, flowers, and stems are used for herbal tea and, in traditional medicine, to treat various ailments, such as cramps, muscle pains, nausea, indigestion, diarrhoea, and infectious diseases (Gulluce et al., 2003).

The aim of this study was to investigate the *in vitro* antioxidant activities of the essential oils and methanol extracts of *S. spicigera* and *S. cuneifolia*. A survey of the literature reveals few reports on the antioxidative activity of water extract of *S. cuneifolia*, and none for *S. spicigera* (Dorman, Bachmayer, Kosar, & Hiltunen, 2004).

2. Materials and methods

2.1. Collection of plant material

The aerial parts (leaves and flowers) of *S. spicigera* (Voucher No. OE 3259) and *S. cuneifolia* (Voucher No. AA 3437) were collected from the Kilickaya district, Yusufeli, Artvin-Turkey and Sogutlugol plateau, Duzici, Osmaniye-Turkey, when flowering (06th September 2004 and 15th July 2004), respectively. The voucher specimens of these plants have been deposited as the herbarium of the Department of Biology, Cumhuriyet University, Sivas-Turkey (CUFH).

2.2. Extraction of the essential oil

The air-dried and finely ground aerial parts of the plants were submitted, for 3 h, to steam distillation, using a Clevenger apparatus (British type) to produce the essential oils in a yield of 2.05% and 1.90% (v/w), based on the dry weight of the samples from *S. spicigera* and *S. cuneifolia*, respectively.

2.3. Gas chromatography (GC) analysis

The essential oil was analyzed using a Hewlett Packard 5890 II GC equipped with a FID detector and HP-5ms capillary column (30 m × 0.25 mm, film thickness 0.25 µm). Injector and detector temperatures were set at 220 and 290 °C, respectively. Oven temperature was raised from 50 to 240 °C by a rate of 3 °C/min. Flow rate of Helium was 1 ml/min. Diluted samples (1/100 in acetone, v/v) of 1.0 µl were injected manually and in the splitless mode. Quantitative data were obtained electronically from FID area percent data without the use of correction factors.

2.4. Gas chromatography/mass spectrometry (GC/MS) analysis

GC/MS analysis of the essential oils was performed under the same conditions with a GC (column, oven temperature, flow rate of the carrier gas) using a Hewlett Pack-

ard 5890 II GC equipped with a Hewlett Packard 5972 mass selective detector in the electron impact mode (70 eV). Injector and MS transfer line temperatures were set at 220 and 290 °C, respectively. The components were identified, based on the comparison of their relative retention time and mass spectra with those of standards and/or NIST98 and Wiley275 library data of the GC-MS system and literature data (Adams, 2001). The results were also confirmed by the comparison of the compound elution orders with their Kovats indices on non-polar phases reported in the literature (Adams, 2001).

2.5. Preparation of the methanol extracts

The air-dried and finely ground samples were extracted by using the method described elsewhere (Sokmen, Jones & Erturk, 1999). Briefly, the samples, weighing about 100 g, were extracted in a Soxhlet apparatus with methanol (MeOH) at 60 °C for 6 h. The extracts were then filtered and concentrated *in vacuo* at 45 °C, yielding a waxy material (15.42 and 14.35% w/w, respectively). Finally, the extracts were lyophilized and kept in the dark at +4 °C until tested.

2.6. Antioxidant activity

2.6.1. DPPH assay

The hydrogen atom or electron donation ability of the corresponding extracts and some pure compounds was measured from the bleaching of purple-coloured methanol solution of DPPH. This spectrophotometric assay uses the stable radical 2,2'-diphenylpicrylhydrazyl (DPPH) as a reagent (Burits & Bucar, 2000; Cuendet, Hostettmann, & Potterat, 1997). Fifty microlitres of various concentrations of the extracts in methanol were added to 5 ml of a 0.004% methanol solution of DPPH. After a 30 min incubation period at room temperature, the absorbance was read against a blank at 517 nm. Inhibition of free radical, DPPH, in percent (I%) was calculated in following way:

$$I\% = (A_{\text{blank}} - A_{\text{sample}}/A_{\text{blank}}) \times 100,$$

where A_{blank} is the absorbance of the control reaction (containing all reagents except the test compound), and A_{sample} is the absorbance of the test compound. Extract concentration providing 50% inhibition (IC_{50}) was calculated from the graph plotting inhibition percentage against extract concentration. Tests were carried out in triplicate.

2.6.2. β -Carotene–linoleic acid assay

In this assay, antioxidant capacity was determined by measuring the inhibition of the volatile organic compounds and the conjugated diene hydroperoxides arising from linoleic acid oxidation (Dapkevicius, Venskutonis, Van Beek, & Linssen, 1998). A stock solution of β -carotene–linoleic acid mixture was prepared as follows 0.5 mg β -carotene

was dissolved in 1 ml of chloroform (HPLC grade); 25 μ l linoleic acid and 200 mg Tween 40 were added. Chloroform was completely evaporated using a vacuum evaporator. Then, 100 ml of distilled water saturated with oxygen (30 min, 100 ml/min.) were added with vigorous shaking. Two thousand five hundred microlitres of this reaction mixture were dispensed to test tubes and 350 μ l portions of the extracts prepared at 2 g l⁻¹ concentrations were added and the emulsion system was incubated for up to 48 h at room temperature. The same procedure was repeated with the synthetic antioxidant, butylated hydroxytoluene (BHT) as positive control, and a blank. After this incubation period, absorbances of the mixtures were measured at 490 nm. Antioxidative capacities of the extracts were compared with those of BHT and blank.

3. Results and discussion

The oils isolated by hydro-distillation from the aerial parts of *S. spicigera* and *S. cuneifolia* were found to be yellow liquids and obtained in yields of 2.05% and 1.90% (v/w), based on dry weights, respectively.

Forty and twenty-nine components were identified in the essential oils of the plants studied. The major components were carvacrol (42.5% and 67.1%), γ -terpinene (21.5% and 15.2%) and *p*-cymene (20.9% and 6.7%), respectively. The chemical composition of the oils can be seen in Table 1.

The oil of *S. spicigera* from Iran is reported by Sefidkon and Jamzad (2004) to contain thymol (35.1%), *p*-cymene (22.1%), γ -terpinene (13.7%) and carvacrol (4.0%). In the study of Tumen and Baser (1996), a variation to the main components thymol (19.56–34.9%), *p*-cymene (7.77–34.1%), γ -terpinene (3.35–19.4%) and carvacrol (1.90–26.1%) was observed, due to the collection of plant material of five different localities of Turkey. In our study, together with *p*-cymene and γ -terpinene, a high amount of carvacrol was found, instead of thymol. This is in accordance with Tanker (1962), who identified carvacrol as the main component in the oil of *S. spicigera*.

The oil of *S. cuneifolia* of Turkish origin is reported to be rich in carvacrol and γ -terpinene (Baydar, Sagdic, Ozkan, & Karadogan, 2004), while samples collected from Croatia were found to contain a low percentage of carvacrol, but were rich in α -pinene, limonene, *p*-cymene, linalool, borneol, and β -cubebene (Milos, Radonic, Bezic, & Dunkic, 2001; Skočibušić & Bezic, 2004). Considerable differences were found to exist in the amounts of these compounds, depending on the locality, the season and the stage of development (Milos et al., 2001).

Our study supports the view that a genuine chemotype for the essential oil of *S. cuneifolia* of Turkish origin is that of carvacrol.

The literature outlines different approaches for determination of the antioxidant activities of the plant extracts. Therefore, different methodological approaches lead to scattered results, which are hardly comparable and often conflicting (Koleva, van Beek, Linssen, de Groot, &

Table 1

Chemical composition of the essential oils of *S. spicigera* and *S. cuneifolia*

No.	K.I. ^a	Components	Composition (%)	
			<i>S. spicigera</i>	<i>S. cuneifolia</i>
1	890	α -Thujene	1.5	0.5
2	898	α -Pinene	1.2	0.8
3	911	Camphene	0.3	0.1
4	935	Sabinene	0.1	–
5	938	β -Pinene	0.1	–
6	945	1-Octene-3-ol	0.6	0.4
7	953	β -Myrcene	2.1	1.8
8	967	α -Phellandrene	0.3	0.2
9	973	<i>p</i> -Mentha-1(7), 8-diene	0.1	0.1
10	979	α -Terpinene	2.1	2.1
11	989	<i>p</i> -Cymene	20.9	6.7
13	1004	Z- β -Ocimene	0.4	–
14	1016	E- β -Ocimene	0.4	0.1
15	1026	γ -Terpinene	21.5	15.2
16	1040	<i>cis</i> -Sabinene hydrate	0.1	tr ^b
17	1062	Terpinolene	0.2	0.2
18	1075	<i>trans</i> -Sabinene hydrate	0.1	–
19	1078	Linalool	–	0.1
22	1155	Borneol	0.2	0.1
24	1168	Terpinen-4-ol	0.2	0.3
25	1187	α -Terpineol	0.1	tr
26	1192	<i>cis</i> -Dihydro carvone	0.1	tr
27	1201	<i>trans</i> -Dihydro carvone	tr	–
29	1246	Carvacrol methyl ether	1.7	0.1
30	1279	Carvone	–	tr
33	1312	Thymol	0.3	0.3
34	1322	Carvacrol	42.5	67.1
38	1399	α -Copaene	0.1	–
39	1411	β -Bourbonene	tr	–
42	1448	β -Caryophyllene	0.6	2.6
44	1459	β -Copaene	tr	–
45	1470	Aromadendrene	tr	0.2
46	1486	α -Caryophyllene	tr	0.1
47	1511	γ -Murolene	0.1	–
48	1516	Germacrene D	0.4	–
50	1531	Viridiflorene	–	0.2
51	1531	Bicyclogermacrene	0.3	–
52	1544	β -Bisabolene	0.4	–
53	1551	γ -Cadinene	0.1	–
54	1560	δ -Cadinene	0.1	tr
55	1580	<i>trans</i> - α -Bisabolene	tr	–
56	1619	Spathulenol	0.2	0.1
57	1624	Caryophyllene oxide	tr	0.1
		Total	99.4	99.5

^a K.I., Kovats index relative to *n*-alkanes on non-polar HP-5ms column.^b Tr, traces \leq 0.06%.

Evstatieva, 2002; Mantle et al., 1998; Ruberto & Baratta, 2000; Zygadlo, Lamarque, Maestri, & Grosso, 1995). A plethora of different antioxidant assays is available and, because results rely on different mechanisms, they strictly depend on the oxidant/antioxidant models employed and on lipophilic/hydrophilic balance (Frankel, Huang, Kanner, & German, 1994). A single/substance/single-assay produces relative results and it is perceived as a reductive approach whenever a phytocomplex is involved. Therefore, antioxidant activities of the plant extracts studied here were determined by two complementary test systems, namely DPPH free radical-scavenging and β -carotene–linoleic acid systems.

Table 2

Free radical-scavenging capacities of the extracts measured in DPPH and β -carotene-linoleic acid assays^a

Extracts	DPPH ($\mu\text{g/ml}$)	β -Carotene-linoleic acid inhibition (%)
<i>Satureja spicigera</i> (essential oil)	127 \pm 1.63	81.7 \pm 1.14
<i>Satureja spicigera</i> (methanol extract)	267 \pm 1.42	65.9 \pm 1.77
<i>Satureja cuneifolia</i> (essential oil)	89.1 \pm 2.29	93.7 \pm 1.83
<i>Satureja cuneifolia</i> (extract)	68.0 \pm 1.76	72.9 \pm 2.16
<i>p</i> -Cymene	NA ^b	NA
γ -Terpinene	NA	NA
Carvacrol	245 \pm 3.26	54.5 \pm 1.42
BHT	18.0 \pm 0.40	96.6 \pm 1.29
Ascorbic acid	3.80 \pm 0.10	94.5 \pm 2.14
Curcumin	7.80 \pm 0.30	89.3 \pm 1.86
α -Tocopherol	6.50 \pm 0.70	96.7 \pm 1.72

^a Results are means of three different experiments.^b NA, Not active.

In both the test systems, all of the extracts and essential oils exhibited remarkable antioxidant activities. In general, samples obtained from *S. cuneifolia* exerted greater antioxidant activities than did those obtained from *S. spicigera*. In general, essential oils showed greater activity than those of the extracts except for *S. cuneifolia* essential oil in the DPPH system. Results of the antioxidant activity test systems can be seen in Table 2.

In the DPPH test system, free radical-scavenging activity of *S. spicigera* oil was determined to be 127 \pm 1.63 $\mu\text{g/ml}$, whereas IC₅₀ value of *S. cuneifolia* was 89.1 \pm 2.29 $\mu\text{g/ml}$. In the second case, antioxidant activities of the oil were 81.7 \pm 1.14% and 93.7 \pm 1.83%, respectively. When compared, methanol extracts of *S. cuneifolia* exhibited greater antioxidant activity than did *S. spicigera* extracts, in both the test systems.

As far as our literature survey could ascertain, methanol extract of *S. cuneifolia* showed better activity than did the aqueous one measured by the same DPPH method (Dorman et al., 2004).


Lamiaceae species are known to produce a diverse array of secondary metabolites, such as volatile and non-volatile terpenes, hydroxybenzoates, hydroxycinnamates and flavonoids, among others. The cardinal mode of action of natural antioxidants is their ability to scavenge free radicals before they can initiate free radical chain reactions in cellular membranes or lipid-rich matrices in foodstuffs, cosmetics or pharmaceutical preparations (Dorman & Hiltunen, 2004).

When compared with the antioxidative potentials of the standard compounds used in this study (BHT, ascorbic acid, curcumin and α -tocopherol), both essential oils and methanolic extracts of the *Satureja* species studied exerted remarkable antioxidant activity.

References

- Adams, R. P. (2001). *Identification of essential oils components by gas chromatography/quadrupole mass spectroscopy*. IL, USA: Allured Publishing Corporation.
- Baydar, H., Sagdic, O., Ozkan, G., & Karadogan, T. (2004). Antibacterial activity and composition of essential oils from *Origanum*, *Thymbra* and *Satureja* species with commercial importance in Turkey. *Food Control*, 15, 169–172.
- Burits, M., & Bucar, F. (2000). Antioxidant activity of *Nigella sativa* essential oil. *Phytotherapy Research*, 14, 323–328.
- Cuendet, M., Hostettmann, K., & Potterat, O. (1997). Iridoid glucosides with free radical-scavenging properties from *Fagraea blumei*. *Helvetica Chimica Acta*, 80, 1144–1152.
- Dapkevicius, A., Venskutonis, R., Van Beek, T. A., & Linssen, P. H. (1998). Antioxidant activity of extracts obtained by different isolation procedures from some aromatic herbs grown in Lithuania. *Journal of the Science of Food and Agriculture*, 77, 140–146.
- Dorman, H. J. D., Bachmayer, O., Kosar, M., & Hiltunen, R. (2004). Antioxidant properties of aqueous extracts from selected Lamiaceae species grown in Turkey. *Journal of Agricultural and Food Chemistry*, 52, 762–770.
- Dorman, H. J. D., & Hiltunen, R. (2004). Fe (III) reductive and free radical-scavenging properties of summer savory (*Satureja hortensis* L.) extract and subfractions. *Food Chemistry*, 88, 193–199.
- Frankel, E. N., Huang, S. W., Kanner, J., & German, J. B. (1994). Interfacial phenomena in the evaluation of antioxidants: bulk oils versus emulsions. *Journal of Agricultural and Food Chemistry*, 42, 1054–1059.
- Gulluce, M., Sokmen, M., Daferera, D., Agar, G., Ozkan, H., Kartal, N., et al. (2003). In vitro antibacterial, antifungal and antioxidant activities of the essential oil, methanol extracts of herbal parts and callus cultures of *Satureja hortensis* L.. *Journal of Agricultural and Food Chemistry*, 51, 3958–3965.
- Koleva, I. I., van Beek, T. A., Linssen, J. P. H., de Groot, A., & Evstatieva, L. N. (2002). Screening plant extracts for antioxidant activity: a comparative study on three testing methods. *Phytochemical Analysis*, 13, 8–17.
- Madsen, H. L., & Bertelsen, G. (1995). Spices as antioxidants. *Trends in Food Science and Technology*, 6, 271–277.
- Mantle, D., Anderton, J. G., Falkous, G., Barnes, M., Jones, P., & Perry, E. K. (1998). Comparison of methods for determination of total antioxidant status: application to analysis of medicinal plant essential oils. *Comparative Biochemistry and Physiology Part B*, 121, 385–391.
- Milos, M., Radonic, A., Bezic, N., & Dunkic, V. (2001). Localities and seasonal variations in the chemical composition of essential oils of *Satureja montana* L. and *S. cuneifolia* ten. *Flavour Fragrance Journal*, 16, 157–160.
- Ruberto, G., & Baratta, M. T. (2000). Antioxidant activity of selected essential oil components in two lipid model systems. *Food Chemistry*, 69, 167–174.

- Sefidkon, F., & Jamzad, Z. (2004). Essential oil composition of *Satureja spicigera* (C. Koch) Boiss from Iran. *Flavour Fragrance Journal*, 19, 571–573.
- Skočibušić, M., & Bezić, N. (2004). Phytochemical analysis and in vitro antimicrobial activity of two *Satureja* species essential oils. *Phytotherapy Research*, 18, 967–970.
- Sokmen, A., Jones, B. M., & Erturk, M. (1999). The in vitro antibacterial activity of Turkish plants. *Journal of Ethnopharmacology*, 67, 79–86.
- Tanker, M. (1962). Studies on *Origanum heracleoticum* L., *majorana onites* (L.) Benth. and *Satureja spicigera* (C. Koch) Boiss used as thyme in Turkey. Dissertation, Istanbul University.
- Tumen, G., & Baser, K. H. C. (1996). Essential oil of *Satureja spicigera* (C. Koch) Boiss from Turkey. *Journal of Essential Oil Research*, 8, 57–58.
- Zygadlo, J. A., Lamarque, A. L., Maestri, D. M., & Grosso, N. R. (1995). Use of essential oils as natural antioxidants. *Grasasy Aceites*, 46, 285–288.



ELSEVIER


<http://www.elsevier.com>

FOOD CHEMISTRY

ISSN: 0308-8146

Abstracting / indexing

- » BIOSIS
- » CAB Abstracts
- » Chemical Abstracts
- » Chemical Engineering
- » Biotechnology Abstracts
- » Current Contents
- » EMBASE
- » EMBiology
- » Food Science and Technology Abstracts
- » Global Health
- » Nutrition Abstracts
- » Publications in Food Microbiology
- » SCISEARCH
- » Science Citation Index
- » Scopus
- » Sociedad Iberoamericana de Informacion Cientifica (SIIC) Data Bases



© Copyright 2007 Elsevier | <http://www.elsevier.com>

FOOD CHEMISTRY

Aims and Scope

Food Chemistry publishes original research papers dealing with the chemistry and biochemistry of foods and raw materials covering the entire food chain from 'farm to fork.' Topics include:

- Chemistry relating to major and minor components of food, their nutritional, physiological, sensory, flavour and microbiological aspects;
- Bioactive constituents of foods, including antioxidants, phytochemicals, and botanicals. Data must accompany sufficient discussion to demonstrate their relevance to food and/or food chemistry;
- Chemical and biochemical composition and structure changes in molecules induced by processing, distribution and domestic conditions;
- Effects of processing on the composition, quality and safety of foods, other bio-based materials, by-products, and processing wastes;
- Chemistry of food additives, contaminants, and other agro-chemicals, together with their metabolism, toxicology and food fate.

Analytical Section

Analytical papers related to the microbiological, sensory, nutritional, physiological, authenticity and origin aspects of food. Papers should be primarily concerned with new or novel methods (especially instrumental or rapid) provided adequate validation is described including sufficient data from real samples to demonstrate robustness. Papers dealing with significant improvements to existing methods, or data from application of existing methods to new foods, or commodities produced in unreported geographical areas, will also be considered.

- Methods for the determination of both major and minor components of food especially nutrients and non-nutrient bioactive compounds (with putative health benefits) will be considered.
- Results of method inter-comparison studies and development of food reference materials for use in the assay of food components;
- Methods concerned with the chemical forms in food, nutrient bioavailability and nutritional status;
- General authentication and origin [e.g. Country of Origin Labelling (COOL), Protected Designation of Origin (PDO), Protected Geographical Indication (PGI), Certificate of Specific Character (CSC)] determination of foods (both geographical and production including commodity substitution, and verification of *organic, biological and ecological* labelling) providing sufficient data from authentic samples should be included to ensure that interpretations are meaningful.

Managing Editor

PROFESSOR G.G. BIRCH

School of Food Biosciences,
University of Reading,
Whiteknights, PO Box 226,
Reading RG6 6AP, UK
e-mail: foodchemedoffice@btopenworld.com

Editor (Analytical Methods)

DR. P.M. FINGLAS

Institute of Food Research,
Norwich Research Park,
Colney, Norwich NR4 7UA, UK
e-mail: paul.finglas@bbsrc.ac.uk

North American Editor

DR. F. SHAHIDI

Dept. of Biochemistry, Memorial University of Newfoundland,
St John's, Newfoundland A1B 3X9, Canada
e-mail: fshahidi@mun.ca

Editor

DR. J.P. ROOZEN

Department of Agrotechnology and Food Sciences,
Wageningen University and Research Center, Biotechnion 172,
PO Box 8129, 6700 EV Wageningen, The Netherlands
e-mail: jacques.roozen@wur.nl

Associate Editors

DR. R.A. FRAZIER

School of Food Biosciences, University of Reading,
Whiteknights, PO Box 226, Reading RG6 6AP, UK
e-mail: r.a.frazier@reading.ac.uk

DR. S. ELMORE

Department of Food Biosciences,
University of Reading, Whiteknights,
PO Box 226, Reading RG6 6AP, UK
e-mail: j.s.elmore@reading.ac.uk

Editorial Board

C. Alasavar

Tubitak Marmara Research Centre,
Food Institute, Turkey

A.T. Andrews

University of Wales Institute, UK

R.C. Berger

Universität Hannover, Germany

T. Beta

University of Manitoba, Canada

P.M. Dey

Royal Holloway, University of London, UK

N.A.M. Eskin

University of Manitoba, Winnipeg, Canada

M.H. Gordon

University of Reading, UK

A.L. Halmos

Department of Food Science, RMIT University,
Melbourne, Australia

M. Jenner

Welcombe, Devon, UK

M.Y. Jung

Department of Food Science and Technology,
Woosuk University, Jeonbuk, Republic of Korea

J.F. Kennedy

University of Birmingham, UK

P. Kilmartin

University of Auckland, New Zealand

J. Lakkis

Pfizer Inc., Morris Plains, NJ, USA

G. Lisinska

Agricultural University, Wroclaw, Poland

I.M. Mackie

Rowett Research Institute, Aberdeen, UK

M. Mathlouthi

University of Reims, France

R.B. Pegg

The University of Georgia, USA

V. Piironen

University of Helsinki, Finland

S. Porretta

Experimental Station for the
Food Preserving Industry, Parma, Italy

P. Puwastien

Institute of Nutrition, Mahidol University
(INMU), Salaya, Phuthamonthon,
Nakhon Pathom, Thailand

E. Risvik

Norwegian Food Research Institute,
Oslo, Norway

R.S. Shallenberger

Cornell University, Geneva, New York, USA

K. Thurlow

LGC Ltd, Teddington, UK

F. Toldrá

Institute of Agrochemistry and Food
Technology (CSIC), Valencia, Spain

R. Tsao (Rong Cao)

Food Research Program, Agriculture and
Agri-Food, Ontario, Canada

R.E. Wrolstad

Oregon State University, USA

V.A. Yaylayan

McGill University, Canada

L. Yu

University of Maryland, USA

J. Zhengyu

Southern Yangtze University, PR China

Analytical Methods

Y. Bao

University of East Anglia, Norwich, UK

L. Castle

Central Science Laboratory, Sand Hutton,
York, UK

A. Ismail

Universiti Putra Malaysia,
Salangor, Malaysia

M. Jagerstad

Department of Food Science,
Swedish University of Agricultural Sciences,
Uppsala, Sweden

J.A. Monro

New Zealand Institute for Crop and
Food Research Ltd, New Zealand

B. Ou

Brunswick Laboratories, Wareham,
Massachusetts, USA

A. Polesello

Istituto Sperimentale per la Valorizzazione
Tecnologica dei Prodotti Agricoli, Milano,
Italy

B. Saad

School of Chemical Sciences, Universiti Sains,
Malaysia


A.J. Tüdös


Shell Global Solutions International BV,
The Netherlands


F. Ulberth


European Commission,
DG Joint Research Centre, Geel, Belgium


Food Chemistry, Volume 100, Issue 1, 2007, Pages 255-263
 Yasunori Hamazu, Frédéric Forest, Kohzy Hiramatsu and Mitsukimi Sugimoto
 SummaryPlus | Full Text + Links | PDF (517 K) | View Related Articles


-  37. ☐


Simultaneous application of transglutaminase and high pressure to improve functional properties of chicken meat gels
Food Chemistry, Volume 100, Issue 1, 2007, Pages 264-272
 Pilar Trespalacios and Reyes Pla
 SummaryPlus | Full Text + Links | PDF (218 K) | View Related Articles
-  38. ☐


Olive oil volatile compounds, flavour development and quality: A critical review
Food Chemistry, Volume 100, Issue 1, 2007, Pages 273-286
 C.M. Kalva, M.S. Allen, D.R. Bedgood, Jr, A.G. Bishop, P.D. Prenzler and K. Robards
 SummaryPlus | Full Text + Links | PDF (306 K) | View Related Articles
-  39. ☐


Combined effect of light salting, modified atmosphere packaging and oregano essential oil on the shelf-life of sea bream (*Sparus aurata*): Biochemical and sensory attributes
Food Chemistry, Volume 100, Issue 1, 2007, Pages 287-296
 Antonios E. Goulas and Michael G. Kontominas
 SummaryPlus | Full Text + Links | PDF (169 K) | View Related Articles
-  40. ☐


Influence of flavour transfer between different gel phases on perceived aroma
Food Chemistry, Volume 100, Issue 1, 2007, Pages 297-305
 Alice B. Nongonierma, Bernard Colas, Mark Springett, Jean-Luc Le Quéré and Andrée Voilley
 SummaryPlus | Full Text + Links | PDF (213 K) | View Related Articles
-  41. ☐


Chymotryptic hydrolysates of α -kafirin, the storage protein of sorghum (*Sorghum bicolor*) exhibited angiotensin converting enzyme inhibitory activity
Food Chemistry, Volume 100, Issue 1, 2007, Pages 306-311
 Vasudeva Kamath, Sajeeda Niketh, Arun Chandrashekar and P.S. Rajini
 SummaryPlus | Full Text + Links | PDF (152 K) | View Related Articles
-  42. ☐


Determination of ochratoxin A in beer by LC-MS/MS ion trap detection
Food Chemistry, Volume 100, Issue 1, 2007, Pages 312-317
 M. Reinsch, A. Töpfer, A. Lehmann, I. Nehls and U. Panne
 SummaryPlus | Full Text + Links | PDF (132 K) | View Related Articles
-  43. ☐

Cadmium-binding protein components of flaxseed: Influence of cultivar and location
Food Chemistry, Volume 100, Issue 1, 2007, Pages 318-325
 B. Dave Oomah, Brian Berekoff, Eunice C.Y. Li-Chan, Giuseppe Mazza, Edward O. Kenaschuk and Scott D. Duguid
 SummaryPlus | Full Text + Links | PDF (131 K) | View Related Articles
-  44. ☐

Effect of carotenoids from red pepper and marigold flower on pigmentation, sensory properties and fatty acid composition of rainbow trout
Food Chemistry, Volume 100, Issue 1, 2007, Pages 326-330
 Yasemen Yanar, Hakan Büyükçapar, Mahmut Yanar and Mustafa Göcer
 SummaryPlus | Full Text + Links | PDF (113 K) | View Related Articles
-  45. ☐

Homo-monoterpenic compounds as chemical markers for *Cydonia oblonga* Miller
Food Chemistry, Volume 100, Issue 1, 2007, Pages 331-338
 Carla Sousa, Branca M. Silva, Paula B. Andrade, Patrícia Valentão, Artur Silva, Federico Ferreres, Rosa M. Seabra and Margarida A. Ferreira
 SummaryPlus | Full Text + Links | PDF (175 K) | View Related Articles
-  46. ☒


The in-vitro antioxidative properties of the essential oils and methanol extracts of *Satureja spicigera* (K. Koch.) Boiss. and *Satureja cuneifolia* ten
Food Chemistry, Volume 100, Issue 1, 2007, Pages 339-343
 Ozgur Eminagaoglu, Bektas Tepe, Onder Yumrutas, H. Askin Akpulat, Dimitra Daferera, Moschos Polissiou and Atalay Sokmen
 SummaryPlus | Full Text + Links | PDF (106 K) | View Related Articles
-  47. ☐

Flavonol glycosides from whole cottonseed by-product
Food Chemistry, Volume 100, Issue 1, 2007, Pages 344-349
 Anna Lisa Piccinelli, Attilio Veneziano, Siro Passi, Francesco De Simone and Luca Rastrelli
 SummaryPlus | Full Text + Links | PDF (154 K) | View Related Articles
-  48. ☐

Prevention of the deterioration of polyphenol-rich beverages
Food Chemistry, Volume 100, Issue 1, 2007, Pages 350-355
 H. Aoshima and S. Ayabe
 SummaryPlus | Full Text + Links | PDF (220 K) | View Related Articles

Thomson.com

THOMSON

 Advanced Search[About Scientific](#)[Scientific Press Room](#)[Contact Scientific](#)


EN : JP : CN : KR : ES : PT

Thomson Scientific : Master Journal List : Journal Search

JOURNAL SEARCH

SCIENCE CITATION INDEX EXPANDED
JOURNAL LIST
Total journals: 6822

[A][B][C][D][E][F][G][H][I][J][K][L][M][N][O][P][Q][R][S][T][U][V][W][X][Y][Z]

"F" Journals 71-80 (of 143) 

FORMAT FOR PRINT A-Z

[Academic](#)
[Corporate](#)
[Engineering & Technology](#)
[Government](#)
[Intellectual Property](#)
[Non-profit](#)
[Pharma & Biotechnology](#)[Product and Customer Support](#)
[Training](#)[Products & Solutions A-Z](#)[Product Login](#)[Free Resources](#)
[Conferences & Events](#)

FOOD AND DRUG LAW JOURNAL

Quarterly

ISSN: 1064-590X

FOOD DRUG LAW INST, 1000 VERMONT AVE NW, SUITE 1200, WASHINGTON, USA, DC, 20005-4903

FOOD AND NUTRITION BULLETIN

Quarterly

ISSN: 0379-5721

INT NUTRITION FOUNDATION, 150 HARRISON AVE, BOSTON, USA, MA, 02111

FOOD AUSTRALIA

Monthly

ISSN: 1032-5298

AUSTRALIAN INST FOOD SCIENCE TECHNOLOGY INC, PO BOX 1303, WATERLOO D C, AUSTRALIA, NSW, 2017

FOOD BIOPHYSICS

Quarterly

ISSN: 1557-1858

SPRINGER, 233 SPRING STREET, NEW YORK, USA, NY, 10013

FOOD BIOTECHNOLOGY

Tri-annual

ISSN: 0890-5436

TAYLOR & FRANCIS INC, 325 CHESTNUT ST, SUITE 800, PHILADELPHIA, USA, PA, 19106

FOOD CHEMISTRY

Monthly

ISSN: 0308-8146

ELSEVIER SCI LTD, THE BOULEVARD, LANGFORD LANE, KIDLINGTON, OXFORD, ENGLAND, OXON, OX5 1GB

FOOD CONTROL

Bimonthly

ISSN: 0956-7135

ELSEVIER SCI LTD, THE BOULEVARD, LANGFORD LANE, KIDLINGTON, OXFORD, ENGLAND, OXON, OX5 1GB

FOOD HYDROCOLLOIDS

Bimonthly

ISSN: 0268-005X

ELSEVIER SCI LTD, THE BOULEVARD, LANGFORD LANE, KIDLINGTON, OXFORD, ENGLAND, OXON, OX5 1GB