

EFFECTS OF LAND-USE REGIME ON SOIL ERODIBILITY INDICES AND SOIL PROPERTIES IN UNYE, TURKEY

Murat Yilmaz^{1*}, Ayhan Usta², Lokman Altun³ and Fahrettin Tilki⁴

¹Faculty of Forestry, Duzce University, 81620 Duzce, Turkey

²Eastern Black Sea Forestry Research Institute, 61300 Trabzon, Turkey

³Faculty of Forestry, Blacksea Technical University, 61080 Trabzon, Turkey

⁴Faculty of Forestry, Artvin Coruh University, 08000 Artvin, Turkey

SUMMARY

We evaluated the effects of land-use regime on soil erodibility indices and several soil properties in forested, deforested, and cultivated areas in the village of Unye, Turkey. Twelve sample plots (spaced 150 m apart) with northern aspects were established in each land-use regime, and samples were taken at soil depths of 0–20, 20–50, and 50–80 cm. Soil organic matter (SOM), soil reaction (pH), total lime (CaCO_3), texture (sand, silt, and clay), dispersion ratio (DR), erosion ratio (ER), colloid-moisture equivalent ratio (C-MER), structural stability index (SSI), field capacity (FC), wilting point (WP), and available water capacity (AWC) were analyzed. The average (of the three soil depths) AWC, FC, and WP values were not affected by the site, although site, soil depth, or both significantly affected other analyzed soil variables. Deforestation and subsequent tillage practices resulted in an almost 20% decrease in clay content, a 33% decrease in SOM, a 15% decrease in AWC, a 51% decrease in total CaCO_3 , a 24% decrease in SSI, a 60% increase in DR, and a 98% increase in ER relative to undisturbed forest soil. At cultivated and forested sites, the ER and DR increased with increasing soil depth. At deforested sites, ER and DR were lowest at 50–80 cm. SOM was the highest at 0–20 cm in the forested sites. Decreasing SOM, clay content, and SSI, as well as increasing DR and ER were outcomes of deforestation. These results indicate that the conversion of forest into cropland deteriorates some soil properties, especially SOM and SSI, and alters the stability of soil aggregates, thus increasing the susceptibility of deforested sites to erosion.

KEYWORDS:

Land use, soil erosion, soil properties, Turkey.

INTRODUCTION

One of the most important issues in Turkish forestry is forest degradation and subsequent forest clearing. The natu-

ral forest ecosystems of the eastern Black Sea region are excessively devastated relative to other forested regions in Turkey. This area of fairly high and steep mountains receives significant precipitation and is intensively populated. The main causes of forest devastation in this region are tillage practices and cultivation (tea or hazelnut) in lower elevation sub-regions (600–1200 m), and transhumance (cultivation of grass and meadow vent, and provision of combustion material) in higher regions of forested areas (1200–1700 m).

Soils in Turkey are increasingly susceptible to erosion owing to the conversion of forests to croplands. Land-use regimes affect soil properties and soil susceptibility to erosion, and deforestation and subsequent tillage practices can change the stability of soil aggregates and deteriorate soil properties [1–5]. According to Knuti et al. [6], agricultural practices in forest areas damage soil quality and increase soil erosion. Boyle [7] and Mroz et al. [8] stated that total tree harvesting may have several negative effects on forest soils, including nutrient removal in the harvested material, increased erosion rates or percolation losses of nutrients, and soil compaction. Likens et al. [9] reported on extensive nutrient losses (particularly of $\text{NO}_3\text{-N}$ and Ca) following deforestation. Conversion of forest and grassland into agricultural land is of considerable concern worldwide in the context of environmental degradation and global climate change [10, 11]. Deforestation also has important consequences for water conservation [2].

The objectives of this study were to examine the effect of land-use regime on soil erodibility indices and other properties, and to evaluate the changes in soil properties resulting from deforestation or cultivation in the village of Unye, Turkey.

MATERIALS AND METHODS

The study was conducted in the village of Unye, on the northeastern coast of Turkey (41°08'–41°05' N, 37°09'–

37°21' E). The study area of about 260 ha extends in the east-west direction, of which 34.6% is deforested which has been clear-cut about 20 years ago and cultivated until now. An additional 30.7% has been under cultivation for many years, and the remainder of the study area is maintained as forest. The elevation of the site ranges from 150 to 650 m above sea level. The landscape in the area faces north, with a slope ranging from 30 to 50%. The region is characterized by a warm and humid Black Sea climate. Mean annual rainfall is approximately 1100 mm, and 60% of which falls in autumn and winter months. The mean annual temperature is 13.4 °C [12].

Fagus orientalis, *Carpinus orientalis*, and *Quercus* spp. are the predominant tree species in the forested area. Fruits and vegetables (e.g., corn, potato, cabbage) and hazelnuts are common crops in the cultivated parts of the study area. As is common in this region, *Fagus orientalis* and *Quercus* spp. were the predominant vegetation in the area prior to deforestation; the vegetation was cleared approximately 20 years ago, and these sites have commonly been used for hazelnut cultivation.

Soil sampling and analyses

Soils were examined from three different land-use regimes (forested, deforested, and cultivated). From each site, 12 samples located 150 m apart were systematically collected for detailed analysis. The samples were taken from three soil depths, 0–20, 20–50, and 50–80 cm, at each site.

Particle size distribution and soil organic carbon were determined using disturbed soil samples passed through a 2-mm sieve using the Bouyoucos hydrometer method [13, 14] and the modified Walkley-Black wet oxidation procedure, respectively. Soil organic matter (SOM) was calculated by multiplying soil organic carbon by 1.72 [15, 16]. Calcium carbonate content (CaCO_3) and soil acidity (pH) were measured using the procedures outlined by Arp [16] and Page et al. [17]. Dispersion ratio (DR) was determined according to the methods described by Middleton [18]. Total clay (TC) and total silt (TS) contents were determined from the particle size distribution using the hydrometer method [19]. The clay and silt fractions obtained by chemical dispersion were taken as TC and TS, while water-dispersible clay and silt (WDCS) was obtained as above, except that no chemical dispersant was used. Colloid-moisture equivalent ratio (C-MER) and erosion ratio (ER) [20], field capacity (FC), wilting point (WP), available water capacity (AWC) [21], and structural stability index (SSI) [22] were also determined.

Statistical analyses

Analyses of variance (ANOVA) were performed to determine the effects of land-use regime and soil depth on the chemical and physical properties of the soil. Arcsine transformation was performed on percentage data, but the means based on the original measurements are presented in the tables. Significant differences between variables were de-

termined by Duncan's new multiple range test. Statistical analyses were performed using SPSS.

RESULTS AND DISCUSSION

ANOVA revealed significant differences among the sites and soil depths, or both, for sand, silt, clay, DR, ER, C-MER, SSI, SOM, pH, and total CaCO_3 ($p < 0.05$) (Tables 1 and 2). FC and WP were not significantly affected by site or soil depth.

The sand, silt, and clay contents averaged across the three soil depths showed significant differences among sites, but there were no significant differences in sand, silt, and clay contents among soil depths at each site (Tables 1 and 2). The average sand content was significantly higher at the deforested site than at the other two sites (67.6%, 73.3%, and 67.1% for the cultivated, deforested, and forested lands, respectively). The average silt content (of all three depths) was highest in the forested soil, and the clay content was higher in the forested and cultivated sites (Table 1).

More clay accumulated at lower depths (20–80 cm) in the cultivated and forested sites. However, the soils at the deforested site had a higher sand content and lower clay content (17.5 to 18.4%) at each depth (Table 2). This may be the result of translocation of finer particles to lower depths (>80 cm), or the movement of finer particles to other areas via erosion, thus leaving the coarser particles in the soils of the deforested site. This scenario is supported by ER and DR results; the average ER and DR values were highest in the deforested site at 0–50 cm. ER and DR increased with soil depth at the cultivated and forested sites, but were lowest in the 50–80 cm soil depth at the deforested site. When averaged over the three soil depths, DR and ER were highest at the deforested site (12.1% and 28.4%, respectively). Soil depth also significantly affected DR and ER. Compared to the soil surface (0–20 cm) at the cultivated and forested sites, the deeper soils had higher DR and ER. The 50–80 cm deep soils at the deforested site had the lowest DR and ER values, 10.1% and 25.0%, respectively (Table 2).

According to Karagül [3], DR and susceptibility to erosion were higher at lower soil depths. In our study, increases in clay content at 50–80 cm depths in the forest and cultivated sites may have caused higher DR at the shallower depths. Nkana and Tonye [4] found that soil sand content was significantly affected by site, and cultivated lands usually have higher sand content in the surface soils than forested sites. As in our study, Hajabbasi et al. [2] found a higher clay content at lower soil depths (30–130 cm) in de-graded and cultivated sites; however, in contrast to our findings, they indicated that clay content was higher in the upper soils (0–30 cm) of a forested site. Some studies have reported higher clay content in cultivated sites [3, 23, 24], whereas Celik [5] found higher clay levels in a forested site. In our study, clay content was higher in the

cultivated and forest sites than in the deforested site, and there was no significant difference between cultivated and forested sites. Decreases in clay content and increases in sand content during the last 25 years at the degraded site in our study could increase the susceptibility of this site to erosion.

Similar to our study, Korkanç [25] and Karagül [3] found a lower DR in forested sites than in cultivated ones. When DR was lower than 10%, soil susceptibility to erosion was low [20, 26]. We found the average DR to be lower than 10% in the forested and cultivated sites, but higher than 10% at all depths in the deforested site. Thus, susceptibility to erosion was higher at the deforested site than at the other two sites. High DR and low C-MER resulted in high ER, as found by Korkanç [25]. High levels of SOM at the forested site resulted in low DR and lower susceptibility to erosion, as reported by Balcı [27] and Karagül [3].

Although site and soil depth did not significantly affect FC and WP, the deeper soils (20–50 and 50–80 cm) had a lower WP (26.1% and 25.4%, respectively) relative to the surface soil (0–20 cm; 30.1%) at the forested site (Table 2). However, WP at the cultivated and deforested sites was lowest in the surface soils (0–20 cm; 27.9% and 25.9%, respectively).

The average FC values (of the three soil depths) were 37.3%, 36.7%, and 35.9% for the cultivated, deforested, and forested sites, respectively. A higher FC was found at the lower depths (20–50 and 50–80 cm) in the cultivated and forested sites, but FC was higher in the surface soils at the deforested site, as with WP.

No significant difference was found between the sites in average AWC (8.26%, 7.20%, and 8.44% for the cultivated, deforested, and forested sites, respectively; Table 1). Soil depth, however, did affect AWC; at the 50–80 cm

TABLE 1 - Effect of land-use regime on soil properties.

Soil property**	Land-use regime	Mean	Standard Deviation	F-ratio	Significance Level
Sand (%)	Cultivated	67.6 b*	9.27	4.85	0.010
	Deforested	73.3 a	9.64		
	Forested	67.1 b	8.90		
Silt (%)	Cultivated	9.5 b	2.97	3.54	0.033
	Deforested	9.0 b	3.26		
	Forested	10.9 a	3.76		
Clay (%)	Cultivated	22.9 a	8.25	3.36	0.038
	Deforested	17.7 b	8.34		
	Forested	22.0 a	10.4		
DR (%)	Cultivated	9.35 b	4.13	6.02	0.003
	Deforested	12.1 a	5.52		
	Forested	7.55 b	6.38		
ER (%)	Cultivated	16.5 b	7.55	10.51	0.000
	Deforested	28.4 a	17.2		
	Forested	14.3 b	15.2		
C-MER	Cultivated	0.60 a	0.14	6.56	0.002
	Deforested	0.48 b	0.30		
	Forested	0.59 a	0.20		
SSI (%)	Cultivated	27.5 a	8.47	7.64	0.001
	Deforested	22.3 b	7.46		
	Forested	29.2 a	7.06		
FC (%)	Cultivated	37.3 a	9.96	0.15	0.861
	Deforested	36.7 a	12.17		
	Forested	35.9 a	6.54		
WP (%)	Cultivated	28.98 a	9.74	0.40	0.672
	Deforested	29.55 a	11.47		
	Forested	27.43 a	7.23		
AWC (%)	Cultivated	8.26 a	3.15	1.26	0.288
	Deforested	7.20 a	3.48		
	Forested	8.44 a	4.14		
SOM (%)	Cultivated	1.78 b	1.24	6.40	0.002
	Deforested	2.08 b	0.88		
	Forested	3.09 a	2.27		
pH	Cultivated	6.72 a	0.71	3.88	0.024
	Deforested	6.35 b	0.70		
	Forested	6.28 b	0.65		
CaCO ₃ (%)	Cultivated	10.1 b	17.78	2.87	0.041
	Deforested	2.59 ab	5.30		
	Forested	5.26 a	14.31		

DR: dispersion ratio, ER: erosion ratio, C-MER: colloid-moisture equivalent ratio, SSI: structural stability index, FC: field capacity, WP: wilting point, AWC: available water capacity, SOM: soil organic matter. *: Means in a row followed by the same letter are not significantly different at $p < 0.05$. **: Values are the averages obtained from three depths (0–20, 20–50 and 50–80 cm).

TABLE 2 - Effect of land-use regime and soil depth on soil properties.

Land-use regime	Depth (cm)	Soil property	Mean	Std. Dev.	F-ratio	Sig. Level	Soil property	Mean	Std. Dev.	F-ratio	Sig. Level
Cultivated	0-20	Sand (%)	69.7 a	10.23	1.34	0.24	SSI (%)	26.4 a	8.41	1.90	0.68
	20-50		67.0 a	8.98				28.7 a	7.73		
	50-80		66.4 a	9.04				27.6 a	9.92		
Deforested	0-20		73.3 a	8.91				22.0 a	6.56		
	20-50		72.3 a	8.94				23.0 a	6.88		
	50-80		74.3 a	11.53				22.0 a	9.26		
Forested	0-20		67.9 a	8.84				29.3 a	7.54		
	20-50		67.4 a	8.14				28.7 a	5.96		
	50-80		65.6 a	10.79				29.8 a	8.48		
Cultivated	0-20	Silt (%)	9.8 a	4.13	1.63	0.13	FC (%)	34.1 a	9.62	0.51	0.84
	20-50		9.1 a	2.61				37.9 a	10.06		
	50-80		9.4 a	1.92				40.0 a	10.16		
Deforested	0-20		9.2 a	2.81				34.3 a	9.17		
	20-50		10.3 a	3.20				37.0 a	11.89		
	50-80		7.3 a	3.31				38.8 a	15.25		
Forested	0-20		11.6 a	3.36				37.5 a	7.06		
	20-50		10.9 a	3.46				34.7 a	6.15		
	50-80		10.4 a	4.95				35.2 a	6.69		
Cultivated	0-20	Clay (%)	20.5 a	8.63	1.06	0.40	WP (%)	27.9 a	10.29	0.67	0.71
	20-50		24.0 a	7.92				29.1 a	9.71		
	50-80		24.2 a	8.44				30.0 a	10.0		
Deforested	0-20		17.5 a	8.20				25.9 a	7.44		
	20-50		17.5 a	8.09				30.1 a	11.5		
	50-80		18.4 a	9.33				32.7 a	14.2		
Forested	0-20		20.5 a	10.67				30.1 a	6.91		
	20-50		21.6 a	9.63				26.1 a	7.16		
	50-80		24.0 a	12.20				25.4 a	7.56		
Cultivated	0-20	DR (%)	8.6 ab	2.18	2.16	0.04	AWC (%)	6.1 a	2.61	1.97	0.58
	20-50		9.4 ab	3.21				8.8 a	2.22		
	50-80		10.2 ab	6.38				10.0 a	3.52		
Deforested	0-20		13.2 a	5.93				8.5 a	4.83		
	20-50		12.8 a	5.80				7.0 a	1.95		
	50-80		10.1 ab	4.70				6.1 a	2.86		
Forested	0-20		5.6 b	4.00				7.4 a	3.63		
	20-50		8.4 ab	7.45				8.5 a	4.26		
	50-80		9.2 ab	7.68				9.8 a	4.75		
Cultivated	0-20	ER (%)	16.0 b	6.42	2.70	0.01	SOM (%)	3.19 b	1.01	9.45	0.00
	20-50		15.6 b	5.91				1.20 e	0.61		
	50-80		18.0 ab	10.43				0.92 bcd	0.49		
Deforested	0-20		29.9 a	18.48				2.72 bcd	0.64		
	20-50		30.4 a	16.79				1.86 de	0.91		
	50-80		25.0 ab	17.41				1.66 de	0.76		
Forested	0-20		12.8 b	12.31				4.48 a	2.91		
	20-50		15.2 b	18.08				2.82 bc	1.37		
	50-80		15.4 b	16.96				1.52 e	0.40		
Cultivated	0-20	C-MER (%)	0.58 abc	0.13	2.15	0.04	pH	6.55 a	0.49	1.16	0.33
	20-50		0.63 ab	0.14				6.75 a	0.80		
	50-80		0.60 abc	0.16				6.90 a	0.84		
Deforested	0-20		0.50 bc	0.13				6.31 a	0.70		
	20-50		0.47 bc	0.14				6.32 a	0.71		
	50-80		0.48 bc	0.15				6.45 a	0.77		
Forested	0-20		0.52 abc	0.19				6.22 a	0.69		
	20-50		0.61 abc	0.18				6.33 a	0.62		
	50-80		0.58 abc	0.13				6.34 a	0.74		
Cultivated	0-20	CaCO ₃ (%)	7.0 a	11.6	0.86	0.55					
	20-50		10.9 a	20.1							
	50-80		12.5 a	21.2							
Deforested	0-20		3.6 a	7.8							
	20-50		2.3 a	4.2							
	50-80		1.8 a	3.2							
Forested	0-20		3.4 a	10.2							
	20-50		6.7 a	17.1							
	50-80		6.0 a	16.7							

DR: dispersion ratio, ER: erosion ratio, C-MER: colloid-moisture equivalent ratio, SSI: structural stability index, FC: field capacity, WP: wilting point, AWC: available water capacity, SOM: soil organic matter. *: Means in a column with the same letter are not significantly different at p<0.05.

depth, the cultivated and forested sites had significantly high AWCs. Relative to the surface soils (0–20 cm; 8.5%) at the deforested site, the deeper soils (20–50 and 50–80 cm) were characterized by lower AWC (7.0% and 6.1% respectively). However, AWC was lowest in the surface soils of the cultivated and forested sites (Table 2). Although there were no significant differences among sites in our study, AWC was higher in the forested area, as found by Karagul [3]. The high SOM content in the forest soil may have increased the soil AWC, as noted in previous studies [28–30].

The average C-MER was lowest in the deforested site (0.48%; Table 1). The C-MER values indicated a difference between land use regimes (Table 2), and deforested sites had the lowest C-MER for all three soil depths. According to Özhan [16], soils with C-MER values lower than 1.5% are susceptible to erosion. In this study, all sites had C-MER values below this threshold, indicating soil susceptibility to erosion.

SSI was significantly affected by site, but not by soil depth (Tables 1 and 2). The deforested site had the lowest average SSI across the three depths (22.3%), and average SSI was found to be 27.5% and 29.2% for the cultivated and forested sites, respectively. A high soil SSI reduces susceptibility to erosion [22]. Doğan and Güçer [31] and Aşkın [32] found that soils with an SSI lower than 40% were more susceptible to erosion.

CaCO₃ was significantly affected by site and soil depth. The average CaCO₃ across the three depths was highest at the cultivated site (10.1%) and lowest at the deforested site (2.59%). High CaCO₃ can increase aggregate size in soil and reduce erosion.

Soil pH was significantly affected by site, and the average pH across the three soil depths was highest at the cultivated site (6.72). Although not significantly affected by soil depth, pH increased in the lower depths at all three sites. The forest at 0–20 cm had the lowest pH (6.22). In a cultivated site, after forest removal, pH usually does not change, and SOM decreases over time [33]. At our cultivated sites, pH was not too low because of liming practices.

SOM was significantly affected by both site and soil depth. The average organic matter content across the three soil depths at the forested site was significantly higher than at the other two sites (3.09%, 2.08%, and 1.78% for the forested, deforested, and cultivated sites, respectively). In most studies, SOM was found to be high in forest soil [15, 34, 35], due to vegetation cover and humus. In our study, the surface soil (0–20 cm) at all sites had the greatest amount of organic matter. The forest site at 0–20 and 20–50 cm had higher amounts of SOM (4.48% and 2.82%, respectively) relative to the same depths at the deforested and cultivated sites (Table 2). At soil depths of 50–80 cm, SOM content was lowest at the cultivated site. Cultivation was associated with significant changes in total nitrogen, and the losses of nitrogen from deforested and cultivated sites appear to be

particularly important. Tillage practices and cultivation caused organic matter content to decrease. Patrick and Smith [36] reported that total tree harvesting results in up to three times greater nutrient removal, including nitrogen removal, compared to conventional logging. In addition to losses from biomass removal, when little vegetation is present to take up nutrients at deforested sites, nutrients can be lost by increased soil nutrient mobilization and leaching [37]. The presence of macroaggregates is usually and positively associated with SOM concentration [38]. Clearly, as explained earlier, cultivation breaks up soil aggregates and exposes previously inaccessible organic matter to microbial attack and accelerated decomposition and mineralization of SOM [39].

The conversion of forest and pastureland into cropland deteriorates soil properties, reduces SOM, and changes the distribution and stability of soil aggregates [36, 37]. Relative to SOM in forest and pasture soils, SOM in cultivated soils was reduced by 44% and 48% in the 0–10 cm layer, and by 48% and 50% in the 10–20 cm layer over 12 years, respectively [5]. Hajabbasi et al. [2] reported similar findings, noting that deforestation and subsequent tillage practices resulted in a nearly 50% decrease in SOM at a soil depth of 0–20 cm over 20 years in the central Zagros Mountains of Iran.

CONCLUSIONS

Each year, hundreds of hectares of land in Turkey are deforested and converted to cropland. We found that some soil properties have been drastically modified by changes in land-use in the village of Unye. Deforestation and subsequent tillage practices changed the distribution and stability of soil aggregates, deteriorated soil properties, and decreased SOM content, SSI, and clay content. Cultivation caused organic matter content to decrease, and reductions in SOM at deforested sites resulted in high DR and ER and low SSI. Thus, deforestation and cultivation increased soil susceptibility to erosion in this region. Because cultivation following deforestation is a common practice in the region, there is a need for a comprehensive soil conservation program to prevent soil erosion problems in susceptible areas.

REFERENCES

- [1] Kalay, H.Z. and Karagül, R. (1992). Ecological degradation, forest destruction, flood and soil erosion in the East Black Sea region. *Ecology Journal* 2: 23–27 (in Turkish).
- [2] Hajabbasi, M.A., Lalalian, A. and Karimzadeh, R. (1997). Deforestation effects on soil physical and chemical properties, Lordegan, Iran. *Plant Soil* 190: 301–308.
- [3] Karagül, R. (1999). Investigations on soil erodibility and some properties of soils under different land use types in Sogutludere Creek Watershed near Trabzon. *Turkish J. Agric. For.* 23: 53–68 (in Turkish).

- [4] Nkana, J.C.V. and Tonye, J. (2003). Assessment of certain soil properties related to different land use systems in Kaya Watershed of the humid forest zone of Cameroon. *Land Degradation & Development*. 14: 57–67.
- [5] Celik, I. (2005). Land-use effects on organic matter and physical properties of soil in a Southern Mediterranean highland of Turkey. *Soil Tillage & Research*. Res. 83: 270–277.
- [6] Knuti, L.L., Williams, D.L. and Hides, J.C. (1979). *Profitable Soil Management*. Prentice Hall, Englewood Cliffs, Prentice Hall, Inc., NJ.
- [7] Boyle, J.R. (1975). Whole tree harvesting impact on site quality. In *Proc. TAPPI Annu. Meet. NY 24–26 February*, pp. 29–32. Technical Association of Pulp and Paper Industries, Atlanta, GA.
- [8] Mroz G.D, Jurgensen M.F. and Frederick D.J. (1985) Soil nutrient changes following whole tree harvesting on tree-northern hardwood sites. *Soil Science Society of America Journal* 49: 1552–1557.
- [9] Likens G.E., Bormann F.H., Johnson N.M., Fisher D.W. and Pierce R.S. (1970) Effects of forest cutting and herbicide treatment on nutrient budgets in the Hubbard Brook watershed-ecosystem. *Ecological Monographs*. 40(1): 23–47.
- [10] Wali, M.K., Evrendilek, F., West, T., Watts, S., Pant, D., Gibbs, H. and McClead, B. (1999). Assessing terrestrial ecosystem sustainability: usefulness of regional carbon and nitrogen models. *Natural Resources*. 35: 20–33.
- [11] Mahtab, F.U. and Karim, Z. (1992). Population and agricultural land use: towards a sustainable food production system in Bangladesh. *Ambio* 2: 50–55.
- [12] Anonymous (1999). Turkish state meteorological service, Ankara, Turkey.
- [13] Bouyoucos, G.J. (1962). Hydrometer method improved for making particle size analyses of soils. *Agronomy Journal*. 54: 464–465.
- [14] Gülçur, F. (1974). Physical and chemical soil analysis methods. Istanbul University Faculty of Forestry Publications, no: 201, Istanbul, Turkey (in Turkish).
- [15] Nelson, D.W. and Sommers, L.E. (1982). Total carbon, organic carbon, and organic matter. In: Page, A.L. (ed.), *Methods of Soil Analysis. Part 2. Chemical and Microbiological Properties*, second ed. Agronomy Monographs, 9. ASA-SSA, Madison, WI, pp. 539–579.
- [16] Arp, P.A. (1999). *Soils for Plant Growth, Field and Laboratory Manual*. Faculty of Forestry and Environmental Management, University of New Brunswick, Canada.
- [17] Page, A.L., Miller, R.H. and Keeney, D.R. (1982). *Methods of Soil Analysis. Part 2. Chemical and Microbiological Properties*, second ed. Agronomy Monographs 9. ASA-SSA, Madison, WI.
- [18] Middleton H.E. (1930). Properties of soils which influence soil erosion. United States Department of Agriculture. Technical Bulletin. No. 178.
- [19] Gee G.W. and Bauder J.W. (1986). Particle-size analysis. In *Methods of Soil Analysis, Part 1*, Klute A. (ed.). American Society of Agronomy and Soil Science Society of America: Madison, WI, pp. 91–100.
- [20] Özhan, S. (2004). Watershed management. Istanbul University Faculty of Forestry, publication no: 482. Istanbul, Turkey (in Turkish).
- [21] Karaöz, M.Ö. (1989). Methods of analysis for some soil chemical properties (pH, carbonate, salinity, organic matter, total nitrogen, available phosphorus). Istanbul University Faculty of Forestry, Series B 39: 64–82 (in Turkish).
- [22] Leo, W.M. (1963). A rapid method for estimating structural stability of soils. *Soil Science*. 96: 342–346
- [23] Tütüdü, Ö.A. (1981). Comparative investigation of soil physical and chemical properties of a *Picea* forest, a beech forest, a grass field and a cornfield in the Hamsikoy region of Trabzon. Karadeniz Technical University Faculty of Forestry Publications, no: 13, Trabzon, Turkey (in Turkish).
- [24] Erol, A. (2004). Effect of hydro-physical soil properties on soil formation factors in Kose Creek Watershed in Gumushane. PhD thesis. Istanbul University Graduate School of Natural & Applied Sciences, Istanbul, Turkey (in Turkish).
- [25] Korkanç S.Y. (2003). Land use problems and recommended solutions in the vicinity of Bartın (A case study of Iskalan Creek Watershed). PhD thesis, Istanbul University Graduate School of Natural & Applied Sciences, Istanbul, Turkey (in Turkish).
- [26] Balci, A.N. (1996). Soil conservation. Istanbul University Faculty of Forestry, publication no: 3947–439. Istanbul, Turkey (in Turkish).
- [27] Balci, A.N. (1973). Effect of parent material and aspect factors on soil properties related to erodibility. Istanbul University Faculty of Forestry, publication no. 195, Istanbul, Turkey (in Turkish).
- [28] Kantarci, M.D. (2000). Soil science. Istanbul University Faculty of Forestry publication no. 3444/387, Istanbul, Turkey (in Turkish).
- [29] Heiskanen, J., and Makitalo, K. (2002). Soil water retention characteristics of Scots pine and Norway spruce forest sites in Finnish Lapland. *Forest Ecology & Management* 162: 137–152.
- [30] Allen, A. and Chapman, D. (2001). Impacts of afforestation on groundwater resources and quality. *Hydrogeology Journal* 9: 390–400.
- [31] Doğan, O. and Güçer, C. (1976). Description of soil erosion and calculation of soil losses using the Universal equation. Soil and Water Resources National Information Centre. Technical publication no. 24, Ankara, Turkey (in Turkish).
- [32] Aşkın, T. (1997). An investigation determining the structural stability and erodibility of Ordu soils. Masters Thesis. Ondokuz Mayıs University Institute of Natural and Applied Sciences, Department of Soil Science. Samsun, Turkey (in Turkish).
- [33] Jaiyeoba, I.A. (2003). Changes in soil properties due to continuous cultivation in Nigerian Semiarid Savannah. *Soil Tillage & Research* 70: 91–98.
- [34] Abbasi, M.K. and Rasool, G. (2005). Effects of different land-use types on soil quality in the hilly area of Rawalakot Azad Jammu and Kashmir. *Acta Agriculturae Scandinavica Section B Soil Plant Science* 55 (3): 221–228.

- [35] Neufeld, H., Resck, D. and Ayarza, M. (2002). Texture and land use effects on soil organic matter in Cerrado Oxisols, Central Brazil. *Geoderma* 107: 151–164.
- [36] Patrick, J.H. and Smith D.W. (1975). Forest management and nutrient cycling in eastern hardwoods. USDA For. Serv. Res. Pap. ME-4. Northeast For. Exp. Stat. Broomal, PA.
- [37] Ross, S.M. (1993). Organic matter in tropical soils: current conditions, concerns and prospects for conservation. *Progress in Physical Geography* 17: 265–305.
- [38] Singh, S. and Singh, J.S. (1996). Water-stable aggregates and associated organic matter in forest, savanna, and cropland soils of a seasonally dry tropical region. *India Biological Fertility Soils* 22: 76–82.
- [39] Boix Fayos, C., Calvo Cases, A., Imeson, A.C. and Soriano Soto, M.D. (2001). Influence of soil properties on the aggregation of some Mediterranean soils and the use of aggregate size and stability as land degradation indicators. *Catena* 44: 47–67.

Received: January 17, 2007

Revised: March 13, 2007; May 07, 2007

Accepted: May 25, 2007

CORRESPONDING AUTHOR

Murat Yilmaz

Faculty of Forestry

Duzce University

81620 Duzce

TURKEY

Phone: +90 0380 5421137-3207

Fax: +90 0380 5421136

E-mail: yilmaz_m2@ibu.edu.tr