



International Soil Congress (ISC) on
"Natural Resource Management
for Sustainable Development"
June 7-10, 2004 Erzurum-Turkey



SOIL RESPIRATION IN YOUNG AND OLD ORIENTAL SPRUCE STANDS AND IN ADJACENT GRASSLANDS IN ARTVIN, TURKEY.

Aydin TUFEKCIOGLU* Temel SARIYILDIZ, Mehmet KUCUK, Sinan GUNER

Department of Forestry, Kafkas University, 08000, Artvin/Turkey,
email: atufekci27@yahoo.com

ABSTRACT

Soil respiration is a good indicator of soil quality. In this study, influence of species type and sampling time on soil respiration in young and old oriental spruce (*Picea orientalis* (L.) Link.) stands without understory and with *Rhododendron ponticum* L. understory and in adjacent grasslands were investigated in Genya Mountain, Artvin, Turkey. Soil respiration was measured approximately monthly from May to November using the soda-lime technique. Mean daily soil respiration across all sites ranged from 0.26-2.66 g C m⁻² d⁻¹. There were significant differences between grasslands and old forest sites, but there was no significant differences between young forest sites and grasslands. Seasonal changes in soil respiration were strongly related to temperature changes. Over all sites, soil temperature and soil moisture together accounted for 75 % of the seasonal variability in soil respiration. Mean soil respiration rates correlated strongly with fine root (<2 mm) biomass (R =0.91, P<0.001), surface (0-15 cm) soil sand content (R =0.71, P<0.05), surface soil silt content (R =-0.69, P<0.05), and subsurface (15-35 cm) soil pH (R=0.60, P<0.05).

Key words: soil biological activity, oriental spruce, grasslands, C cycle

INTRODUCTION

Soil respiration is the release of CO₂ from soils to atmosphere and it is an important process in carbon cycle in forest and grassland ecosystems. Soils are major global sources and sinks of CO₂ and therefore play an important role in regulating atmospheric concentrations of CO₂. Almost 10% of the atmosphere's CO₂ passes through soils each year. This is more than 10 times the CO₂ released from fossil fuel combustion (Raich and Schlesinger, 1992).

Soil respiration is a sensitive indicator of several essential ecosystem processes, including: metabolic activity in soil, persistence and decomposition of plant residue in soil, and conversion of soil organic carbon to atmospheric CO₂ (Rochette et al., 1997). In addition, Parkin et al. (1996) stated that soil respiration is a good indicator of soil quality.

Soil moisture and soil temperature are two significant determinants of soil respiration (Singh and Gupta, 1977; Raich and Tufekcioglu, 2000). Rochette et al. (1997) observed that soil respiration in moist soil was two to three times greater than in drier soils. Most researchers also reported an increase in soil respiration with increasing soil temperature (e.g., Kowalenko et al., 1978).

Soil respiration varies with vegetation type (Raich and Tufekcioglu, 2000). However, analyzing published soil respiration data Raich and Tufekcioglu (2000) found no



predictable significant ($P < 0.05$) differences in soil respiration between cropped and vegetation-free soils, between grassland and cropped soils or between forested and cropped soils. They did find higher rates of soil respiration in grasslands than in forests, and in grasslands and forests than in adjacent croplands.

Estimates of soil respiration have been made in a variety of ecosystems and have been summarized in reviews by Schlesinger (1977), Singh and Gupta (1977), Raich and Schlesinger (1992) and Raich and Tufekcioglu (2000). Despite this considerable body of information on soil respiration in different parts of the world, there is no soil respiration study in forest and grassland ecosystems of Turkey.

The objectives of this study were to compare rates of soil respiration among young oriental spruce stands (SSY), old oriental spruce stands with no understory (SSO), old oriental spruce stands with *Rhododendron ponticum* L. understory (SSR) and in adjacent grasslands; and to identify the underlying environmental variables most likely causing differences in soil respiration among sites, and among seasons within sites. We hypothesized that grasslands have higher rates of soil biological activity, and therefore have higher rates of soil respiration than do adjacent spruce stands.

MATERIALS AND METHODS

The study site is located at Genya Mountain of Artvin, Turkey. The site with a northern aspect and gentle slope (10-20%), ranges in elevation from 1490 m to 1510 m. Soil at the site is a podsollic well-drained sandy-loam. Soil respiration were measured in young oriental spruce stands (SSY), in old spruce stands with no understory (SSO), in old oriental spruce stands with *Rhododendron ponticum* L. understory (SSR) and in adjacent grasslands. Young stands were around 15 years-old and established after a clearcutting by planting and natural regeneration. Old stands were around 90 years-old with normal canopy cover. Plot sizes were 20 x 20 m. Dominant grass species in the grassland sites were smooth brome (*Bromus inermis* Leysser.), *Agrostis tenuis* L., timothy (*Phleum pratense* L.), Kentucky bluegrass (*Poa pratensis* L.) and *Festuca* sp. Grassland sites were under heavy grazing until last year when grazing was stopped.

Soil samples were taken randomly from 0-15 cm and 15-35 cm soil depths by digging a soil pit in each plot. Soil samples were air-dried, ground and pass through 2 mm mesh-sized sieve. Organic matter contents of the soils were determined according to wet digestion method described by Kalra and Maynard (1991)(Modified Walkley-Balck Method). Soil texture determined by Bouyoucos' Hydrometer Method. Soil pH was determined by a combination glass-electrode in H_2O (soil-solution ratio 1: 2.5).

The biomass of fine (0-2 mm) roots was assessed by collecting six 35-cm deep, 6.4-cm diameter cores per plot in October (Harris et al., 1977; Tufekcioglu et al., 1999). Roots were separated from the soil by soaking in water and then gently washing them over a series of sieves with mesh sizes of 2.0 and 0.5 mm. Roots were sorted into diameter classes of 0-2 mm (fine root), 2-5 mm (small root) and 5-10 mm (coarse root) root classes. The roots from each size category were oven-dried at 65 °C for 24 h and weighed.



Soil respiration rates were measured approximately monthly in three randomly selected locations in each of the three plots per sites from May 2003 to November 2003 using the soda-lime method (Edwards, 1982; Raich et al., 1990). The soda-lime method may underestimate actual soil respiration rates at high flux rates (e.g. Ewel et al., 1987; Haynes and Gower, 1995). However, the method does distinguish between higher and lower flux rates and, therefore, it is an appropriate method for comparing sites.

Buckets 20 cm tall and 27.5 cm in diameter were used as measurement chambers. One day prior to measurements, plastic rings with the same diameter were placed over the soil and carefully pushed about 1 cm into the soil. All live plants inside the plastic rings were cut to prevent aboveground plant respiration. Carbon dioxide was absorbed with 60 g of soda-lime contained in 7.8 cm diameter by 5.1 cm tall cylindrical tins. In the field, the plastic rings were removed, measurement chambers were placed over the tins of soda-lime, and the chambers were held tightly against the soil with rocks. After 24 h the tins were removed, oven dried at 105° C for 24 h, and weighed. Blanks were used to account for carbon dioxide absorption during handling and drying (Raich et al. 1990). Soda-lime weight gain was multiplied by 1.69 to account for water loss (Grogan, 1998). Soil temperature was measured at 5 cm soil depth adjacent to each chamber in the morning. Diurnal variations in soil temperature were expected to be smaller at these sites because of shading of sunlight by the plant canopy. Gravimetric soil moisture was determined by taking soil samples at 0-5 cm depth and drying them at 105 °C for 24 h on the day that the soda-lime tins were removed from the plots.

Statistical comparisons were made using SPSS program. We used ANOVA to compare soil respiration rates, soil temperatures, and soil moisture contents among sites. Paired comparisons among sites and sampling dates were determined with Least Significant Difference test at $\alpha=0.05$. Step-wise multiple regression analysis was performed to evaluate the importance of soil temperature and soil moisture on seasonal soil respiration rates. The possible effects of soil properties and fine root biomass on soil respiration rates were evaluated with correlation analysis.

RESULTS AND DISCUSSION

Mean daily soil respiration ranged from 0.26 to 2.66 g C m⁻² d⁻¹ among all sites (Fig. 1A). These values are within the ranges reported by Singh and Gupta (1977), Hudgens & Yavitt (1997), Coleman (1973), Jurik et al. (1991), Kucera & Kirkham (1971) and Lessard et al. (1994), Tufekcioglu et al. (2001). Highest rates were observed in mid August when soil temperatures were high, while lowest rates were observed in May when soil temperatures were minimal (Fig. 1B). Soil respiration increased from spring to summer and decreased from summer to fall, as is typical in temperate latitudes (e.g., Hudgens and Yavitt, 1997; Kowalenko et al., 1978). Our results indicated that temperature was limiting during the fall and spring and moisture was limiting during the summer or fall. Kowalenko et al. (1978) reported that temperature was limiting during the winter and spring and moisture was limiting during the summer or fall on soil respiration in field soils in Canada.

Soil respiration varied significantly among sites ($P<0.01$). Soil respiration was significantly lower in old spruce stands (SSO and SSR) than in grassland sites (Table 1).



**International Soil Congress (ISC) on
"Natural Resource Management
for Sustainable Development"
June 7-10, 2004 Erzurum-Turkey**



Similar results were reported by Raich and Tufekcioglu (1999). They reported that grasslands had ~20% higher soil respiration rates than did comparable forest stands. Higher soil respiration rates in grassland sites were probably due to higher soil temperatures and fine root biomass values in these sites. Soil temperature and fine root biomass are significant determinants of soil respiration in temperate latitudes (e.g. Tufekcioglu et al., 2001; Kelting et al., 1998).




Figure 1: Mean monthly (± 1 SE) soil respiration rates (A) and soil temperature (B) in young (SSY) and old oriental spruce stands with (SSR) / without understory (SSO) and in adjacent grasslands.

There were significant differences in soil respiration among sampling dates, but with different temporal patterns for forest and grassland vegetation types. Soil respiration increased from May to August in all sites and it decreased from August to November in grasslands while decreasing from August to September, and increasing from September to November in the other sites. These differences in temporal patterns of soil respiration were probably driven by moisture and temperature differences among sites. There was no change in soil moisture from August to September in grassland sites but in the same period, soil moisture content decreased with an average value of 26.3% in the other sites. Transpiration from the aboveground tissues and interception of rain by the forest canopy might account for these soil moisture differences among sites.

Soil temperature varied significantly among sites and sampling dates ($P < 0.01$) (Fig. 1B). Soil temperatures in the grassland sites were significantly different from those in the other sites ($P < 0.05$) (Table 1). There was no other significant temperature



**International Soil Congress (ISC) on
"Natural Resource Management
for Sustainable Development"
June 7-10, 2004 Erzurum-Turkey**



differences among sites. Averaged over all sites, soil temperatures were significantly higher in July than in October.

Soil moisture content differed significantly between sampling dates. Overall soil moisture content were significantly higher in October than in May, July and September ($P < 0.05$) (Fig. 2). Mean soil moisture contents (average of five sampling dates) were 35.3%, 31.4%, 35.1% and 27.1 in grasslands, in SSO, in SSY and in SSR sites, respectively (Table 1). Higher soil moisture contents in July, September and October in grassland sites compared to SSO and SSR sites were probably result of higher canopy interception in the forest stands than in grasslands. Cepel (1971) reported that interception ranged from 17-31% for forest stands and 6-17 for grasslands in Turkey.

Within sites, seasonal changes in soil respiration were correlated most highly with soil temperature. When all sites were considered together, mean daily soil respiration varied with soil temperature and moisture ($r^2 = 0.75$, $P < 0.001$):

$$SR = 0.14 T + 0.0429 M - 2.194$$

where SR is the soil respiration rate ($\text{g C m}^{-2} \text{d}^{-1}$), T is morning surface-soil (0-5 cm depth) temperature ($^{\circ}\text{C}$) and M is surface-soil (0-5 cm depth) gravimetric moisture content (% H_2O). All three parameters were significant ($P < 0.01$). According to stepwise regression results, 46% and 29% of variation in soil respiration explained by soil temperature and soil moisture, respectively. This indicated that soil temperature was the most important determinant of soil respiration in these high elevation sites.

Among sites, mean annual soil respiration rate correlated positively with fine root biomass, surface soil (0-15 cm) sand content, mean soil temperature, mean soil moisture and subsurface soil (15-35 cm) pH; and correlated negatively with surface soil silt content ($P < 0.05$) (Table 2). Soil temperature and soil sand content were positively correlated, suggesting that high soil temperatures were associated with sandy soils (Kantarci, 2000).

Soil respiration had the highest correlation with mean fine root biomass ($r = 0.91$). Respiration by roots and their associated microbial components represent a significant part of soil respiration in most ecosystems (Bowden et al., 1993; Kelting et al., 1998). While live roots directly contribute to soil respiration, dead roots and root exudates provide carbon as an energy source and nutrients for microbial biomass. Grayston et al. (1996) reported that root exudates stimulate microbial growth and activity because they are readily assimilated, and they may act as primers for the degradation of existing soil organic matter.

In a native prairie, belowground litter contributed 20-25 %, root respiration contributed 25-30 %, and decay of organic matter contributed 30-35 % of the total soil respiration (Buyanovsky et al., 1987). Root respiration accounted for 33-50 % of total soil respiration in broad-leaved forests, 17-40 % in grasslands, and 12-38 % in crop fields in temperate regions (Raich and Tufekcioglu, 1999).




Figure 2: Mean monthly (± 1 SE) soil moisture contents (0-5 cm depth) in young (SSY) and old oriental spruce stands with (SSR) / without understory (SSO) and in adjacent grasslands

CONCLUSIONS

In this study, grasslands had higher rates of soil respiration than adjacent old forest sites did. These higher rates of soil respiration are evidence of high rates of the biological activity and C cycling through the soil. Our results suggest that old forests might be better than grasslands in terms of carbon accumulation into soil and forest floor in these high elevation sites.

Table 1: Mean soil respiration, soil temperature, soil moisture, soil organic matter, soil sand, clay and silt content, pH and root biomass in the four sites investigated in this study (n=3 plots per site). Root data refer to the surface 35 cm of soil.

Vegetation types	Grassland	Old spruce stand with	Old spruce stand no	Young spruce stand	Old spruce stand with
Mean soil respiration ($\text{g C m}^{-2} \text{d}^{-1}$)	1.68	0.89		1.13	0.59
Mean soil temperature($^{\circ}\text{C}$)	16.6	11.7		13.1	12.5
Mean soil moisture (%)	35.3	31.4		35.1	27.1
Mean soil organic matter (%) 0-15 cm	5.58	7.57		6.54	6.11
15-35 cm	2.90	4.43		3.39	3.69
Mean sand content (%) 0-15 cm	66.4	57.5		58.2	52.3
15-35 cm	44.9	53.24		56.2	51.59
Mean clay content (%) 0-15 cm	16.8	25.5		24.0	23.2
15-35 cm	30.7	27.0		28.4	27.6
Mean silt content (%) 0-15 cm	16.9	17.1		17.8	24.5
15-35 cm	24.4	19.8		15.6	20.8
Mean soil pH 0-15 cm	5.33	5.29		5.41	4.98
15-35 cm	5.57	5.32		5.17	4.83
Mean fine root biomass (<2mm)(gm^{-2})	785	498		564	404



International Soil Congress (ISC) on
"Natural Resource Management
for Sustainable Development"
June 7-10, 2004 Erzurum-Turkey



Table 2: Pearson correlation coefficients among measured variables in the study area (n=12). Asterisks refer the level of significance; *, P<.05; **, P<.01; ***, P<.001

Variables	FRB	T	M	pH	Sand(%)	Clay(%)	Silt(%)	SOC
SR ¹	0.91***	0.83***	0.65*	0.28	0.71*	-0.50	-0.68*	-0.33
FRB ²	1.0	0.87***	0.75**	0.28	0.65*	-0.53	-0.54	-0.23
T ³		1.0	0.43	0.17	0.62*	-0.69*	-0.25	-0.47
M ³			1.0	0.24	0.33	-0.06	-0.54*	0.07
pH				1.0	0.61*	-0.51	-0.49	0.25
SOC ¹					-0.28	0.41	0.01	1.0


SR: soil respiration ($\text{g C m}^{-2} \text{d}^{-1}$); SOC: Mean soil organic matter content (%).

FRB: fine root biomass (kg ha^{-1}),

M: Mean soil moisture (%), T: Mean soil temperature (C°)

REFERENCES

- Bowden, R. D., K.J. Nadelhoffer, R.D. Boone, J.M. Melillo and J.B. Garrison. 1993. Contributions of aboveground litter, belowground litter, and root respiration to total soil respiration in a temperate mixed hardwood forest. *Can.J. For. Res.* 23,1402-1407.
- Buyanovsky, G. A., C.L. Kucera and G.H. Wagner. 1987. Comparative analyses of carbon dynamics in native and cultivated ecosystems. *Ecology* 68 (6), 2023-2031.
- Coleman, D. C. 1973. Soil carbon balance in a successional grassland. *Oikos* 24, 195-199.
- Cepel, N. 1971. Toprak yuzeyine varan yagis miktarina bitkilerin yaptigi etki ve Belgrad Ormaninda yapilan bir arastirmaya ait bes yillik sonuclar. *I.U. Orman fakultesi Dergisi*, XXI, 2/B, s. 54-59.
- Davidson, E. A., E. Belk and R.D. Boone. 1998. Soil water content and temperature as independent or confounded factors controlling soil respiration in a temperate mixed hardwood forest. *Global Change Biology* 4, 217-227.
- Edwards, N. T. 1982. The use of soda-lime for measuring respiration rates in terrestrial systems. *Pedobiologia* 23, 321-330.
- Ewel, K. C., W. C. Cropper and H. L. Gholz. 1987. Soil CO₂ evolution in Florida slash pine plantations. I. Changes through time. *Canadian J. of Forest Research* 17, 325-329.
- Grayston, S. J., D. Vaughan and D. Jones. 1996. Rhizosphere carbon flow in trees, in comparison with annual plants: the importance of root exudation and its impact on microbial activity and nutrient availability. *Applied Soil Ecology* 5, 29-56.
- Grogan, P. 1998. CO₂ flux measurement using soda-lime: Correction for water formed during CO₂ adsorption. *Ecology* 79, 1467-1468.
- Harris, W.F., Kinerson, R.S. Jr and N. T. Edwards. 1977. Comparison of belowground biomass of natural deciduous forest and Loblolly pine plantations. P. 27-29, in *The Belowground Ecosystem: A Synthesis of Plant -Associated Processes*, JK Marshall (Ed.). Range Science Department, Science Series No: 26, Fort Collins, Colorado, USA.
- Haynes, B. E. and S. T. Gower. 1995. Belowground carbon allocation in unfertilized and fertilized red pine plantations in northern Wisconsin. *Tree Physiol.* 15, 317-325.
- Hudgens, D. E. and J. B. Yavitt. 1997. Land-use effects on soil methane and carbon dioxide fluxes in forest near Ithaca, New York. *Ecoscience* 4(2), 214-222.



**International Soil Congress (ISC) on
"Natural Resource Management
for Sustainable Development"
June 7-10, 2004 Erzurum-Turkey**



- Jurik, T. W., G. M. Briggs and D. M. Gates. 1991. Soil respiration of five aspen stands in Northern Lower Michigan. *Am. Midl. Nat.* 126, 68-75.
- Kalra, Y. P. and D. G. Maynard. 1991. *Methods Manual for Forest Soil and Plant Analysis*. Forestry Canada, Northern Forestry Publications. Alberta, Canada.
- Kantarci, M. D. 2000. *Toprak Ilmi*. I.U. Orman Fakultesi Yayinlari, Yayin No: 462. Istanbul, Turkiye.
- Kelting, D. L., J.A. Burger and G.S. Edwards. 1998. Estimating root respiration, microbial respiration in the rhizosphere, and root-free soil respiration in forest soils. *Soil Biology and Biochemistry* 30, 961-968.
- Kowalenko, C. G., K. C. Ivarson and D.R. Cameron. 1978. Effect of moisture content, temperature and nitrogen fertilization on carbon dioxide evolution from field soils. *Soil Biology and Biochemistry* 10, 417-423.
- Kucera, C. L. and D.R. Kirkham. 1971. Soil respiration studies in tallgrass prairie in Missouri. *Ecology* 52, 912-915.
- Lessard, R., P. Rochette, E. Topp, E. Pattey, R.L. Desjardins and G. Beaumon. 1994. Methane and carbon dioxide fluxes from poorly drained adjacent cultivated and forest sites. *Canadian Journal of Soil Science* 74, 139-146.
- Parkin, T. B., J. W. Doran and E. Franco-Wizcaino. 1996. Field and Laboratory tests of soil respiration in *Methods for Assessing Soil Quality*. Ed. J.W. Doran and A.J. Jones. Soil Science Society of America special publication no 49, Madison, Wisconsin, USA.
- Raich, J. W. and W. H. Schlesinger. 1992. The global carbon dioxide flux in soil respiration and its relationship to vegetation and climate. *Tellus* 44B, 81-99.
- Raich, J. W. and A. Tufekcioglu. 2000. Vegetation and soil respiration: Correlations and controls. *Biogeochemistry* 48 (1), 71-90.
- Raich, J. W., R. D. Bowden and P. A. Steudler. 1990. Comparison of two static chamber techniques for determining carbon dioxide efflux from forest soils. *Soil Science Society of America Journal* 54, 1754-1757.
- Rochette, P., B. Ellert, E. G. Gregorich, R. L. Desjardins, E. R. Pattey Lessard and B. G. Johnson. 1997. Description of a dynamic closed chamber for measuring soil respiration and its comparison with other techniques. *Canadian Journal of Soil Science* 77, 195-203.
- Schlesinger, W. H. 1977. Carbon balance in terrestrial detritus. *Ann. Rev. Ecol. Syst.* 8, 51-81.
- Singh, J. S. and S. R. Gupta. 1977. Plant decomposition and soil respiration in terrestrial ecosystems. *Bot. Rev.* 43, 449-528.
- Tufekcioglu, A., J. W. Raich, T. M. Isenhardt and R. C. Schultz. 1999. Fine root dynamics, coarse root biomass, root distribution, and soil respiration in a multi species riparian buffer in Central Iowa, USA. *Agroforestry Systems* 44, 163-174.
- Tufekcioglu, A. J. W. Raich, T. M. Isenhardt and R. C. Schultz. 2001. Soil respiration within riparian buffers and adjacent crop fields. *Plant and Soil* 229: 117-124.
- Tufekcioglu, A. J. W. Raich, T. M. Isenhardt and R. C. Schultz. 2003. Biomass, carbon and nitrogen dynamics of multi-species riparian buffers within an agricultural watershed in Iowa, USA. *Agroforestry Systems* 57: 187-198.